

BIANCHI TYPE II STIFF FLUID MODE IN PRESENCE OF BULK VISCOSITY

B.L. Meena

Department of Mathematics, Govt. P.G. College, Tonk (Rajasthan)

Email: drblmina27@yahoo.co.in

Abstract: Bianchi Type II stiff fluid model in presence of bulk viscosity is investigated. To get the deterministic model of the universe, we assumed that shear (σ) is Proportional to expansion (θ) which leads to $R = S^n$ where R and S are metric potentials and n is a constant. We find that bulk viscosity prevents the matter density to vanish. The spatial volume increases exponentially representing inflationary scenario. The model represents decelerating and accelerating phases of universe which matches with the result of Astronomical observations.

Key Words: Bianchi II, Stiff fluid, Bulk viscosity

2010 Mathematics Subject classification : 83 Cxx, 83 Fxx

1. Introduction

It has been argued for long time that dissipative process may well account for the high degree of isotropy, we observe today (Misner [1]). Dissipative effects including bulk viscosity play a significant role in the study of early evolution of universe. Eckart [2] first developed the relativistic theory of non-equilibrium thermodynamics to study the effect of viscosity. Padmanabhan and Chitre [3] have discussed that the presence of bulk viscosity leads to inflationary like solution in general relativity. The effect of bulk viscosity on cosmological evolution has been investigated by many authors viz. Sahni and Starobinsky [4], Mak and Harko [5,6], Peebles [7], Saha [8], Hu and Meng [9], Ren and Meng [10], Singh et al. [11], Bali and Singh [12], Gagnon and Legoulrgues [13].

2. Metric and Field Equations

We consider Bianchi Type II metric in the form

$$ds^2 = -dt^2 + R^2(dx^2 + dz^2) + S^2(dy - x dz)^2 \quad (1)$$

where R and S are functions of t alone.

Energy momentum tensor in presence of bulk viscosity is given by

$$T_i^j = (\rho + p)v_i v^j + pg_i^j - \zeta\theta (g_i^j + v_i v^j) \quad (2)$$

with

$$g_{ij} v^i v^j = -1 \quad (3)$$

We assume the coordinates to be comoving so that $v^1 = 0 = v^2 = v^3$, $v^4 = 1$ where ρ the matter density, p the isotropic pressure, ζ the coefficient of bulk viscosity and θ the expansion in the model, v^i the flow vector satisfying (3).

Now

$$T_1^1 = p - \zeta\theta, T_2^2 = p - \zeta\theta, T_3^3 = p - \zeta\theta, T_4^4 = -\rho \quad (4)$$

The Einstein field equations

$$R_i^j - \frac{1}{2}R g_i^j = -8\pi T_i^j \quad (5)$$

(In geometrized units $G = 1$, $c = 1$ and taking $\Lambda = 0$)

for the line element (1) leads to

$$\frac{R_{44}}{R} + \frac{S_{44}}{S} + \frac{R_4 S_4}{RS} + \frac{S^2}{4R^4} = -8\pi(p - k) \quad (6)$$

$$\frac{2R_{44}}{R} + \frac{R_4^2}{R^2} - \frac{3S^2}{4R^4} = -8\pi(p - k) \quad (7)$$

$$\frac{2R_4 S_4}{RS} + \frac{R_4^2}{R^2} - \frac{S^2}{4R^4} = -8\pi\rho \quad (8)$$

where subscript 4 denotes differentiation with respect to t and

we assume

$$\zeta\theta = k \quad (9)$$

3. Solution of Field Equations

Equations (6) and (8) lead to

$$\frac{R_{44}}{R} + \frac{S_{44}}{S} + \frac{3R_4 S_4}{RS} + \frac{R_4^2}{R^2} = -8\pi(\rho - p - k) \quad (10)$$

To get determinate solution, we assume

$$(i) \quad \text{stiff fluid condition } \rho = p \quad (11)$$

$$(ii) \quad \sigma \propto \theta \text{ which leads to } R = S^n \quad (12)$$

Now after using (11) and (12) in (10), We have

$$(n+1) \frac{S_{44}}{S} + (2n^2 + 3n) \frac{S^2}{S^2} = 8\pi k \quad (13)$$

which leads to

$$2S_{44} + 4n \frac{S^2}{S} = \frac{16\pi k}{n+1} S \quad (14)$$

Let $S_4 = f(S)$

$$\therefore S_{44} = f f', f' = df/dS$$

$$\text{I.F.} = S^{4n}$$

Solution is

$$\begin{aligned} f^2 S^{4n} &= \frac{16\pi k}{n+1} \int S^{4n+1} dS \\ &= \frac{16\pi k}{(n+1)(4n+2)} S^{4n+2} + \alpha \end{aligned}$$

where α is constant of integration. Thus

$$f^2 = \frac{8\pi k}{(n+1)(2n+1)} S^2 + \alpha S^{-4n} \quad (15)$$

which leads to

$$\frac{dS}{\sqrt{\beta S^2 + \alpha S^{-4n}}} = dt \quad (16)$$

where

$$\beta = \frac{8\pi k}{(n+1)(2n+1)} \quad (17)$$

To get determinate solution, we assume

$$n=1/2 \quad (18)$$

Using (17), (18) in (16), we have

$$\frac{S dS}{\sqrt{(S^2)^2 + \gamma^2}} = \sqrt{\beta} dt \quad (19)$$

where

$$\frac{\alpha}{\beta} = \gamma^2$$

Let

$$S^2 = \tau$$

$$S dS = \frac{d\tau}{2} \quad (20)$$

Equation (19) leads to

$$\sinh^{-1} \left(\frac{\tau}{\gamma} \right) = \sqrt{\beta} t + b$$

which leads to

$$\tau = \gamma \sinh (at+b)$$

Thus we have

$$S^2 = \gamma \sinh (at + b) \quad (21)$$

where

$$a = \sqrt{\beta}$$

Therefore

$$R^2 = S = \sqrt{\gamma} \sinh^{1/2}(at + b) \quad (22)$$

The metric (1) leads to

$$ds^2 = -dt^2 + \sqrt{\gamma} \sinh^{1/2}(at + b) (dx^2 + dz^2) \\ + \gamma \sinh (at + b) (dy - xdz)^2 \quad (23)$$

4. Some Physical and Geometrical Features

We have

$$S^2 = \gamma \sinh(at + b)$$

$$\therefore \frac{S_4}{S} = \frac{a}{2} \coth(at + b)$$

The matter density from equation (8) is given by

$$\begin{aligned} -8\pi\rho &= \frac{2R_4 S_4}{RS} + \frac{R_4^2}{R^2} - \frac{S^2}{4R^4} \\ &= 2 \frac{1}{2} \frac{S_4^2}{S^2} + \frac{1}{4} \frac{S_4^2}{S^2} - \frac{1}{4} \\ &= \frac{5}{4} \frac{S_4^2}{S^2} - \frac{1}{4} \\ &= \frac{5}{4} \frac{a^2}{4} \coth^2(at + b) - \frac{1}{4} \\ 8\pi\rho &= -\frac{5a^2}{16} \coth^2(at + b) + \frac{1}{4} \\ &= 8\pi p \end{aligned} \tag{24}$$

The expansion (θ), the shear (σ), the spatial volume (V^3), the deceleration parameter (q) are given by

$$\begin{aligned} \theta &= \frac{R_4}{R} + \frac{S_4}{S} \\ &= \frac{3}{2} \frac{S_4}{S} \\ &= \frac{3a}{2} \coth(at + b) \end{aligned} \tag{25}$$

$$\begin{aligned}\sigma &= \frac{1}{\sqrt{3}} \left| \frac{R_4}{R} - \frac{S_4}{S} \right| \\ &= \frac{a}{4\sqrt{3}} \coth(at + b)\end{aligned}\quad (26)$$

$$V^3 = R^2 S = S^2 = \gamma \sinh(at + b) \quad (27)$$

$$q = -\frac{\ddot{V}/V}{\dot{V}^2/V^2}$$

Now

$$V^3 = \gamma \sinh(at + b)$$

$$\therefore \frac{\dot{V}}{V} = \frac{a}{3} \coth(at + b)$$

$$\frac{\ddot{V}}{V} - \frac{\dot{V}^2}{V^2} = -\frac{a^2}{3} \operatorname{cosech}^2(at + b)$$

$$\begin{aligned}\frac{\ddot{V}}{V} &= -\frac{a^2}{3} \operatorname{cosech}^2(at + b) + \frac{a^3}{9} \coth^2(at + b) \\ &= -\frac{a^2}{3} [\coth^2(at + b) - 1] + \frac{a^2}{9} \coth^2(at + b) \\ &= \frac{a^2}{3} - \frac{2a^2}{9} \coth^2(at + b)\end{aligned}$$

$$\begin{aligned}\therefore q &= -\frac{9 \left[\frac{a^2}{3} - \frac{2a^2}{9} \coth^2(at + b) \right]}{a^2 \coth^2(at + b)} \\ &= -3 \tanh^2(at + b) + 2\end{aligned}\quad (28)$$

Particle horizon

$$\begin{aligned}
 &= \int_{t_0}^t \frac{dt}{V^3(t)} \\
 &= \int_{t_0}^t \frac{dt}{\gamma \sinh(at + b)} \\
 &= \text{finite}
 \end{aligned} \tag{29}$$

5. Discussion and Conclusion

The reality condition $\rho > 0$ leads to

$$\coth^2(at + b) < \frac{4}{5a^2}$$

The equation (24) indicates that bulk viscosity prevents the matter density to vanish. The model (23) starts with a big-bang at $t = -b/a$ and the expansion decreases with time. Since $\frac{\sigma}{\theta} \neq 0$, hence anisotropy is maintained throughout. The spatial volume increases exponentially with time. Thus the model represents inflationary scenario. The model also represents accelerating and decelerating phases of universe matching with recent astronomical observations. Also the particle horizon is finite, hence the particles are in communicable region. The model in general represents realistic model of the universe.

Acknowledgement

I am extremely thankful to Prof. Raj Bali, CSIR Emeritus Scientist for helpful discussion and suggestion.

References

- [1] Misner, C.W. (1968). The isotropy of the Universe, *Astrophys. J.* 151, 431-457.
- [2] Eckart, C. (1940). The thermodynamics of Irreversible process III Relativistic Theory of the simple fluid, *Phys. Rev.* 58, 919.
- [3] Padmanabhan, T. and Chitre, S.M. (1987). *Phys. Lett. A. Viscous Universe*, 120, 433-436.
- [4] Sahni, V. and Starobinsky, A. (2000). The case for a positive cosmological - Term, *Int. J. Mod. Phys. D9*, 373-443.
- [5] Mak, M.K. and Harko, T. *Europhys. (2001). Exact dissipative cosmologies with stiff fluid, Europhys. Lett.* 56, 762-767.
- [6] Mak, M.K. and Hark, T. (2004). Full Causal dissipative cosmologies with stiff matter, *Int. J. Mod. Phys. D* 13, 273.

- [7] Peebles, P.J.E. (2003). The cosmological and dark energy, *Rev. Mod. Phys.* 75, 559-606.
- [8] Saha, B. (2005). Bianchi type I Universe with viscous fluid, *Mod. Phys. Lett. A*, 20, 2127-2143.
- [9] Hu, M.G. and Meng, X.H. (2006). Bulk viscous cosmology: State finder and entropy, *Phys. Lett. B*, 635, 186-194.
- [10] Ren, J. and Meng, X.H. (2006). Cosmological model with viscosity media (dark energy) described by a effective equation of state, *Phys. Lett. B*, 633, 1-8.
- [11] Singh, C.P., Kumar, S. and Pradhan, A. (2007). Early viscous universe with variable gravitational and cosmological constant. *Class. Quant. Gravity* 24, 455-474.
- [12] Bali, R. and Singh, J.P. (2008). Bulk viscous Bianchi Type I cosmological models with Time-Dependent Cosmological Term, *Int. J. Theor. Phys.*, 47, 3288-3297.
- [13] Gagon, J.S. and Lergourgues, J.J. (2011). Bulk viscosity as an alternative to dark energy, *Cosmology Astropart. Phys.* 1109, 026.