

BIANCHI TYPE VI₀ DARK ENERGY MODELS WITH CONSTANT DECELERATION PARAMETER IN SELF CREATION COSMOLOGY

Vimal Chand Jain

Department of Mathematics, Govt. Engineering College, Ajmer, India
vcjeca@gmail.com

Nikhil Jain

Department of Mathematics, Govt. Women Engg. College, Ajmer, India
nikhiljain@gweca.ac.in

Abstract: We have constructed dark energy cosmological model in an anisotropic Bianchi type VI₀ space time with a variable equation of state (EoS) parameter ω in Barber's second self-creation theory of gravitation. The models are obtained using the special law of variation of Hubble's parameter that yields a constant value of deceleration parameter. Models have been derived using above conditions. Physical and geometrical properties have also been discussed.

Keywords: Bianchi, Dark Energy, Self Creation

Mathematics Subject Classification: 83F05, 85A40

1. Introduction

Recent observations on expansion phenomenon of the universe indicate that the universe is currently experiencing a phase of accelerated expansion. This was first observed from high red shift supernova Ia (Reiss et al. [20,21], Perlmutter et al. [18], Astier et al. [6], Spergel et al.[26] etc.) and same was confirmed by cross checks from the cosmic microwave background radiation (Bennett et al. [11], Abazajian et al.[1-3], Hawkins et al. [15] etc.). The current accelerating expansion of the universe attributed to the fact that our universe is dominated by an unknown dark energy (DE) an exotic energy with negative pressure. The simplest DE candidate is the vacuum energy density which is mathematically equivalent to the cosmological constant Λ . As per Copeland et al. [13] "fine tuning" and the cosmic "coincidence" are the two well-known difficulties of the cosmological constant problems. There are several alternative theories for the dynamical DE scenario which have been proposed by scientists to interpret the accelerating universe. Wang and Tegmark [28] have shown that the universe is actually undergoing an acceleration with repulsive gravity of some strange energy-form i.e. DE at work. DE is a mysterious substance with negative pressure and accounts for nearly 70% of total matter-energy of universe, but has no clear explanation. Karami et al. [16] introduced a

polytropic gas model of DE as an alternative model to explain the accelerated expansion of the universe. Gupta and Pradhan [14] proposed a new candidate known as cosmological nuclear-energy as a possible candidate for the DE.

Bianchi types I-IX cosmological models are important in the study because these are homogeneous and anisotropic from which the process of isotropization of the universe is studied through the passage of time. Moreover, from the theoretical point of view anisotropic universe have a greater generality than FRW isotropic models. The simplicity of the field equations made Bianchi space-times useful in constructing models of spatially homogeneous and anisotropic cosmologies. Considerable works have been done in obtaining various Bianchi type cosmological models and their inhomogeneous generalization. Bianchi type-VI₀ space-time is of special interest in anisotropic cosmology. Barrow [10] pointed out that Bianchi type-VI₀ models of the universe give a better explanation of some of the cosmological problems like primordial helium abundance and they also isotropize in a special sense. Looking to the importance of Bianchi type-VI₀ universes, many authors [7, 8, 19, 22, 27] have studied it in different context. Shri Ram[24, 25] has presented Bianchi type-VI₀ cosmological models filled with dust and perfect fluid in modified Brans-Dicke theory respectively. Adhav et al. [5] studied Bianchi type-VI₀ cosmological models with anisotropic dark energy. Abdussattar and Prajapati [4] obtained a class of bouncing non-singular FRW models by constraining the deceleration parameter (DP) in the presence of an interacting DE represented by a time-varying cosmological constant. They have also discussed the role of deceleration parameter and interacting DE in singularity avoidance. Bisabr [12] has shown that an accelerating expansion is possible in a spatially flat universe for large values of the Brans-Dicke parameter consistent with the local gravity experiments. Yadav and Saha [29] studied DE models with variable equation of state (EoS) parameter. Recently, Saha and Yadav [23] presented a general relativistic cosmological model with time-dependent DP in LRS Bianchitype-II space-time which can be described by isotopic and variable EoS parameter. In this paper, We present general relativistic cosmological models with constant and time-dependent DP in Bianchi type-VI₀ space-time which can be described by isotropic constant and variable EoS parameters. This paper is organized as follows: We present the metric and field equations in Sect.2. In Sect.3, we obtain the solutions of the field equations representing Bianchi type-VI₀ cosmological models with perfect fluid by imposing the condition that the shear scalar is proportional to expansion scalar. We also discuss the physical and geometrical behaviors of the cosmological models with DE in section 4. Concluding remarks are given in Sect.5.

2. Metric and Field Equations

We consider Bianchi VI₀ metric in the form

$$ds^2 = -dt^2 + A^2(t)dx^2 + e^{-2qx} B^2(t)dy^2 + e^{2qx} C^2(t)dz^2 \quad (1)$$

where A, B and C are cosmic-scale functions and q is non-zero constant.

The Einstein's field equations in Barber's [9] second theory of self creation are given by

$$R_i^j - \frac{1}{2} R g_i^j = -8\pi\phi^{-1} T_i^j \quad (2)$$

$$\phi_{;k}^k = \frac{8\pi}{3} \eta T' \quad (3)$$

where $\phi_{;k}^k$ is invariant d'Alembertian and the contracted tensor T' is trace of energy momentum tensor describing all non-gravitational and non-scalar field matter and energy. Here η is a coupling constant to be determined from experiments. Because of the homogeneity condition imposed by the metric, the scalar field ϕ will be a function of t only.

The simplest generalization of the EoS parameter of a perfect fluid may be to determine the EoS parameter separately on each spatial axis by presenting the diagonal form of the energy momentum tensor.

Thus the energy momentum tensor T_i^j is given by

$$T_j^i = \text{diag} [T_1^1, T_2^2, T_3^3, T_4^4]$$

It can be parameterized as

$$T_j^i = \text{diag} [-p_x, -p_y, -p_z, \rho]$$

$$T_j^i = \text{diag} [-\omega_x, -\omega_y, -\omega_z, 1] \rho$$

$$T_j^i = \text{diag} [-\omega, -(\omega + \delta), -(\omega + \gamma), 1] \rho \quad (4)$$

Where ρ is energy density of the fluid. p_x , p_y and p_z are the pressures and ω_x , ω_y and ω_z are the directional EoS parameters along x, y and z axes respectively. We have parameterized the deviation from isotropy by setting $\omega_x = \omega$, where ω is deviation free EoS parameter of the fluid and then introducing skewness parameter. δ and γ are deviation from ω along y and z axes respectively.

In a comoving coordinate system, field equations (2) and (3) for the metric (1) in case of (4), yield the following system of equations

$$\frac{B_{44}}{B} + \frac{C_{44}}{C} + \frac{B_4 C_4}{BC} + \frac{q^2}{A^2} = \frac{8\pi\rho\omega}{\phi} \quad (5)$$

$$\frac{C_{44}}{C} + \frac{A_{44}}{A} + \frac{A_4 C_4}{AC} - \frac{q^2}{A^2} = \frac{8\pi\rho(\omega + \delta)}{\phi} \quad (6)$$

$$\frac{A_{44}}{A} + \frac{B_{44}}{B} + \frac{A_4 B_4}{AB} - \frac{q^2}{A^2} = \frac{8\pi\rho(\omega + \gamma)}{\phi} \quad (7)$$

$$\frac{A_4 B_4}{AB} + \frac{B_4 C_4}{BC} + \frac{A_4 C_4}{AC} - \frac{q^2}{A^2} = -\frac{8\pi\rho}{\phi} \quad (8)$$

$$\frac{B_4}{B} - \frac{C_4}{C} = 0 \quad (9)$$

$$\phi_{44} + \phi_4 \left(\frac{A_4}{A} + \frac{B_4}{B} + \frac{C_4}{C} \right) = \frac{8\pi}{3} \lambda (3\omega + \delta + \gamma - 1) \rho \quad (10)$$

where the sub-indices 4 denotes ordinary differentiation with respect to t .

3. Solutions of Field Equations

Integrating (9) we obtain

$$C = lB \quad (11)$$

where l is constant of integration.

From equations (6) and (11), we obtain

$$\frac{B_{44}}{B} + \frac{A_{44}}{A} + \frac{B_4 A_4}{BA} - \frac{q^2}{A^2} = \frac{8\pi\rho(\omega + \delta)}{\phi} \quad (12)$$

From equations (7) and (12), we have

$$\delta = \gamma \quad (13)$$

Hence skewness in y and z directions is equal.

In view of equation (11), equations (5) to (10) reduce to

$$2\frac{B_{44}}{B} + \frac{B_4^2}{B^2} + \frac{q^2}{A^2} = \frac{8\pi\rho\omega}{\phi} \quad (14)$$

$$\frac{A_{44}}{A} + \frac{B_{44}}{B} + \frac{A_4 B_4}{AB} - \frac{q^2}{A^2} = \frac{8\pi\rho(\omega + \gamma)}{\phi} \quad (15)$$

$$2\frac{A_4 B_4}{AB} + \frac{B_4^2}{B^2} - \frac{q^2}{A^2} = \frac{8\pi\rho}{\phi} \quad (16)$$

$$\phi_{44} + \phi_4 \left(\frac{A_4}{A} + 2 \frac{B_4}{B} \right) = \frac{8\pi}{3} \lambda (3\omega + 2\gamma - 1) \rho \quad (17)$$

Equations (14) to (17) are four independent equations in A , B , ρ , ω , ϕ and γ (six unknowns), hence we need two more conditions to solve these equations.

- (i) Firstly, we apply the law of variation for Hubble parameter, which yields a constant value of deceleration parameter

$$H = kR^{-n}$$

Which yields

$$H = k(ABC)^{-n/3} \quad (18)$$

Where $k > 0$ and $n \geq 0$ are constants

From equation (18), we have

$$\frac{1}{3} \left(\frac{A_4}{A} + \frac{B_4}{B} + \frac{C_4}{C} \right) = k(ABC)^{-n/3} \quad (19)$$

which on integration gives

$$ABC = (k_1 t + k_2)^{3/n} \quad \text{for } n \neq 0 \quad (20)$$

where k_1 and k_2 are constant of integration.

The deceleration parameter so obtained is

$$q = n - 1 \quad (21)$$

- (ii) Secondly we assume expansion(θ) is proportional to shear(σ), which leads to $A = \beta B^m$ (22)

Where β is constant of integration.

From equations (11), (20) and (22) we get

$$A = \left(\frac{\beta}{l} \right)^{m/m+1} (k_1 t + k_2)^{3m/n(m+1)} \quad (23)$$

$$B = \left(\frac{1}{\beta l} \right)^{1/m+1} (k_1 t + k_2)^{3/n(m+1)} \quad (24)$$

$$C = l \left(\frac{1}{\beta l} \right)^{1/m+1} (k_1 t + k_2)^{3/n(m+1)} \quad (25)$$

Where k_1 and k_2 are constant of integration. Using equations (23) – (25), metric (1) takes the form

$$ds^2 = -dt^2 + \left(\frac{\beta}{l} \right)^{2m/m+1} (k_1 t + k_2)^{6m/n(m+1)} dx^2 + e^{-2qx} \left(\frac{1}{\beta l} \right)^{2/m+1} (k_1 t + k_2)^{6/n(m+1)} dy^2 + e^{2qx} l^2 \left(\frac{1}{\beta l} \right)^{2/m+1} (k_1 t + k_2)^{6/n(m+1)} dz^2 \quad (26)$$

using suitable transformations we can obtain the metric in following form

$$ds^2 = -\frac{dT^2}{k_1^2} + \left(\frac{\beta}{l} \right)^{2m/m+1} T^{6m/n(m+1)} dX^2 + e^{-2qX} \left(\frac{1}{\beta l} \right)^{2/m+1} T^{6/n(m+1)} dY^2 + e^{2qX} l^2 \left(\frac{1}{\beta l} \right)^{2/m+1} T^{6/n(m+1)} dZ^2 \quad (27)$$

where $t = k_1 t + k_2$, $X = x$, $Y = y$, $Z = z$

4. Physical and Geometrical Properties

Scalar expansion (θ) is given as

$$\theta = 3H \quad (28)$$

For our model (27), (θ) is given as

$$\theta = \frac{3k_1 (m+2)}{n(m+1)T} \quad (29)$$

Components of Hubble Parameter are obtained as

$$H_1 = \frac{A_4}{A} = \frac{3k_1 m}{n(m+1)T} \quad (30)$$

$$H_2 = \frac{B_4}{B} = \frac{3k_1}{n(m+1)T} \quad (31)$$

$$H_3 = \frac{C_4}{C} = \frac{3k_1}{n(m+1)T} \quad (32)$$

and Hubble parameter is obtained as

$$H = \frac{k_1(m+2)}{n(m+1)T} \quad (33)$$

Shear (σ^2) is given as

$$\sigma^2 = \frac{1}{4} \left[\frac{9k_1^2 m(m-4)}{n^2(m+1)^2 T^2} \right] \quad (34)$$

Anisotropy parameter is obtained as

$$A_m = \frac{1}{3} \left[\sum_{i=1}^3 \left(\frac{\Delta H_i}{H} \right)^2 \right] \quad (35)$$

For the model (27) it is given as

$$A_m = \frac{1}{27} (3m^2 - 8m + 6) \quad (36)$$

From equation (16) we have

$$\frac{\rho}{\phi} = -\frac{1}{8\pi} \left[\frac{18k_1^2 m}{n^2(m+1)^2 T^2} + \frac{9k_1^2}{n^2(m+1)^2 T^2} - \frac{q^2}{\left(\frac{\beta}{l}\right)^{2m/m+1} T^{6m/n(m+1)}} \right] \quad (37)$$

From equation (14), we get

$$\omega = - \left[\frac{\frac{6k_1^2(3-nm-n)}{n^2(m+1)^2 T^2} + \frac{9k_1^2}{n^2(m+1)^2 T^2} + \frac{q^2}{\left(\frac{\beta}{l}\right)^{2m/m+1} T^{6m/n(m+1)}}}{\frac{18k_1^2 m}{n^2(m+1)^2 T^2} + \frac{9k_1^2}{n^2(m+1)^2 T^2} - \frac{q^2}{\left(\frac{\beta}{l}\right)^{2m/m+1} T^{6m/n(m+1)}}} \right] \quad (38)$$

Equation (15) gives

$$\gamma = \frac{\left[\frac{6k_1^2(3-nm-n)}{n^2(m+1)^2 T^2} - \frac{3k_1^2 m(n+nm-9m)}{n^2(m+1)^2 T^2} + \frac{9k_1^2 m}{n^2(m+1)^2 T^2} - \frac{q^2}{\left(\frac{\beta}{l}\right)^{2m/m+1} T^{6m/n(m+1)}} \right]}{\left[\frac{18k_1^2 m}{n^2(m+1)^2 T^2} + \frac{9k_1^2}{n^2(m+1)^2 T^2} - \frac{q^2}{\left(\frac{\beta}{l}\right)^{2m/m+1} T^{6m/n(m+1)}} \right]} \quad (39)$$

From equation (13) we get

$$\delta = \frac{\left[\frac{6k_1^2(3-nm-n)}{n^2(m+1)^2 T^2} - \frac{3k_1^2 m(n+nm-9m)}{n^2(m+1)^2 T^2} + \frac{9k_1^2 m}{n^2(m+1)^2 T^2} - \frac{q^2}{\left(\frac{\beta}{l}\right)^{2m/m+1} T^{6m/n(m+1)}} \right]}{\left[\frac{18k_1^2 m}{n^2(m+1)^2 T^2} + \frac{9k_1^2}{n^2(m+1)^2 T^2} - \frac{q^2}{\left(\frac{\beta}{l}\right)^{2m/m+1} T^{6m/n(m+1)}} \right]} \quad (40)$$

From equations (17), (38), (39) and using the condition

$$n = \frac{6m}{2(2m+1)}, \text{ we get}$$

$$T^2 \phi_{44} + T p_1 \phi_4 - p_2 \phi = 0 \quad (41)$$

Where

$$p_1 = \frac{3(m+2)k_1}{n(m+1)} \text{ and}$$

$$p_2 = \frac{3k_1^2(3-nm-n)}{n^2(m+1)^2} - \frac{3k_1^2 m(n+nm-9m)}{n^2(m+1)^2} + \frac{18k_1^2 m}{n^2(m+1)^2} + \frac{9k_1^2}{n^2(m+1)^2} - \frac{q^2}{\left(\frac{\beta}{l}\right)^{2m/m+1}}$$

On solving equation (41) we get

$$\phi = c_1 T^{\frac{1}{2}} \left(-i\sqrt{p_1} \sqrt{\frac{(p_1-1)^2}{p_2} - 4 - p_1 + 1} \right) + c_2 T^{\frac{1}{2}} \left(i\sqrt{p_1} \sqrt{\frac{(p_1-1)^2}{p_2} - 4 - p_1 + 1} \right) \quad (42)$$

Where c_1 and c_2 are constants of integration.

Energy density (ρ) is given as

$$\rho = -\frac{1}{8\pi} \left[\frac{18k_1^2 m}{n^2 (m+1)^2 T^2} + \frac{9k_1^2}{n^2 (m+1)^2 T^2} - \frac{q^2}{\left(\frac{\beta}{l}\right)^{2m/m+1} T^{6m/n(m+1)}} \right] \times$$

$$\left[c_1 T^{\frac{1}{2} \left(-i\sqrt{p_1} \sqrt{\frac{(p_1-1)^2}{p_2} - 4 - p_1 + 1} \right)} + c_2 T^{\frac{1}{2} \left(i\sqrt{p_1} \sqrt{\frac{(p_1-1)^2}{p_2} - 4 - p_1 + 1} \right)} \right] \quad (43)$$

The spatial volume (V) is evaluated as

$$V = R^3 = ABC = \frac{1}{(\beta l)^{\frac{m}{3(m+1)}}} T^{\frac{m+2}{n(m+1)}} \quad (44)$$

Hence we can observe that $V \rightarrow 0$ as $T \rightarrow 0$ and $V \rightarrow \infty$ as $T \rightarrow \infty$ with $\frac{m+2}{n(m+1)} > 0$ but $V \rightarrow \infty$ as $T \rightarrow 0$ and $V \rightarrow 0$ as $T \rightarrow \infty$ with $\frac{m+2}{n(m+1)} < 0$.

Now

$$\frac{\sigma}{\theta} = \frac{1}{2} \left[\frac{m\sqrt{m-4}}{m+2} \right] \quad (45)$$

Jerk Parameter $\{j(t)\}$ for the model (27) is given as

$$j(t) = \frac{1}{H^3} \frac{\ddot{R}}{R} = \frac{[(m+2) - n(m+1)][(m+2) - 2n(m+1)]n(m+1)}{(m+2)^3} T^{3-3n(m+1)} \quad (46)$$

5. Conclusion

In this paper we have presented Bianchi type VI₀ dark energy models with constant deceleration parameter in self creation cosmology. The model obtained in this paper is of great interest and may be useful in self-creation cosmology to study the large scale dynamics of physical space.

It is also observed that when $T \rightarrow 0$ the physical quantities for the model (27) like scalar expansion (θ), shear scalar (σ) and Hubble parameter (H) diverge for $m \neq -1, n \neq 0$. Thus the model (27) starts with a big-bang at $T=0$. The scalar field (ϕ) is given by equation (42). we can observe that $V \rightarrow 0$ as $T \rightarrow 0$ and $V \rightarrow \infty$ as $T \rightarrow \infty$ with $\frac{m+2}{n(m+1)} > 0$ but $V \rightarrow \infty$ as $T \rightarrow 0$ and $V \rightarrow 0$ as $T \rightarrow \infty$ with $\frac{m+2}{n(m+1)} < 0$.

Anisotropy Parameter is obtained is constant for the model (27). The model (27) has a POINT TYPE singularity at $T=0$ (MacCallum [17]) when $m \neq -1, n \neq 0$. Jerk parameter is given by equation (46).

Acknowledgement: The authors are thankful to the Referee for valuable comments and suggestions.

References

- [1] Abazajian, K., et al. (2003). "The First Data Release of the Sloan Digital Sky Survey", *Astron. J.* 126, 2081.
- [2] Abazajian, K., et al. (2004). *arXiv: astro.ph/0410239*.
- [3] Abazajian, K., et al. (2004). "The Second Data Release of the Sloan Digital Sky Survey", *Astron. J.* 128, 502.
- [4] Abdussattar, Prajapati, S.R. (2011). "Role of deceleration parameter and interacting dark energy in singularity avoidance", *Astrophys. Space Sci.* 331, 657.
- [5] Adhav, K.S., et al. (2011). "Bianchi type-VI₀ cosmological models with anisotropic dark energy", *Astrophys. Space Sci.* 332, 497.
- [6] Astier, P., et al. (2006). "The Supernova Legacy Survey: measurement of, and w from the first year data set", *Astron. Astrophys.* 447, 31.
- [7] Bali, R., Banerjee, R., Banerjee, S.K. (2008). "Bianchi type VI₀ magnetized bulk viscous massive string cosmological model in General Relativity", *Astrophys. Space Sci.* 317, 21.
- [8] Bali, R., Pradhan, A., Hassan, A. (2008). "Bianchi Type VI₀ Magnetized Barotropic Bulk Viscous Fluid Massive String Universe in General Relativity", *Int. J. Theor. Phys.* 47, 2594.
- [9] Barber, G.A. (1982). "On two 'self creation' Cosmologies", *Gen. Relativity Gravitation* 14, 117.
- [10] Barrow, J.D. (1984). "Helium formation in cosmologies with anisotropic curvature", *Mon. Not. Astron. Soc.* 211, 221.
- [11] Bennet, C.L., et al. (2003). "First year Wilkinson microwave anisotropy probe (WMAP)", *Astrophys. J. Suppl.* 148, 1
- [12] Bisabr, Y.: *Gen. Relative. Gravit.* DOI 10. 1007/s 10714-011-1281-8.
- [13] Copeland, E.J., Sami, M., Tsujikawa, S. (2006). "Dynamics of Dark Energy", *Int. J. Mod. Phys.D* 15, 1753.

- [14] Gupta, R.C., Pradhan, A., "Genesis of Dark Energy: Dark Energy as Consequence of Release and Two-Stage Tracking of Cosmological Nuclear Energy", *Int. J. Theor. Phys.* doi:10, 1007/10773-010-0261-1
- [15] Hawkins, E., et al. (2003). "The 2dF Galaxy Redshift Survey: correlation functions, peculiar velocities and the matter density of the Universe", *Mon. Not. R. Astron. Soc.* 346, 78.
- [16] Karami, et al. (2009). "Interacting polytropic gas model of phantom dark energy in non-flat universe", *Eur. Phys. J.C* 64, 85.
- [17] MacCallum, M.A.H. (1971). "A class of homogeneous cosmological models III: Asymptotic behaviour", *Comm. Math. Phys.* **20**, 57.
- [18] Perlmutter, S., et al. (1997). "Measurements* of the Cosmological Parameters Ω and Λ from the First Seven Supernovae at $z \geq 0.35$ ", *Astrophys. J.* 483, 565.
- [19] Pradhan, A., Bali, R. (2008). "Magnetized Bianchi Type VI_0 Barotropic Massive String Universe with Decaying Vacuum Energy Density Λ ", *EJTP* 19, 91.
- [20] Riess, A.G., et al. (1998). "Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant", *Astron. J.* 116, 1009.
- [21] Riess, A.G., et al. (2004). "Type Ia Supernova Discoveries at $z > 1$ from the Hubble Space Telescope: Evidence for Past Deceleration and Constraints on Dark Energy Evolution*", *Astrophys. J.* 607, 665.
- [22] Roy, S.R., Singh, J.P. (1983). *ActaPhysiacAustriaca* 55, 57.
- [23] Saha, B., Yadav, A.K., "Dark energy model with variable q and ω in LRS Bianchi-II space-time", *Astrophys. Space Sci.* DOI 10.1007/s/10509- 012-1070-1.
- [24] Shri Ram (1986). *J. Math. Phys.* 27, 660.
- [25] Shri Ram, J. (1985). "Bianchi type- VI_0 solutions in modified Brans–Dicke cosmology", *Math. Phys.* 26, 2916.
- [26] Spergel, D.N., et al. (2003). "First-Year Wilkinson Microwave Anisotropy Probe (WMAP)* Observations: Determination of Cosmological Parameters", *Astrophys. J. Suppl.* 148, 175.
- [27] Tikekar, R., Patel, L.K. (1994). "Some exact solutions in Bianchi VI_0 string cosmology", *Pramana-J. Phys.* 42, 483.
- [28] Wang, Y., Tegmark, M. (2004). "Gravitational clustering of relic neutrinos and implications for their detection", *Phys. Rev. Lett.* 92, 241301.
- [29] Yadav, A.K., Saha, B. (2012). "LRS Bianchi-I anisotropic cosmological model with dominance of dark energy", *Astrophys. Space Sci.* 337, 759